


Fast radio bursts: a new exotic puzzle in astrophysics

SERGEI POPOV (SAI MSU)

CO-AUTHORS: K. POSTNOV, M. PSHIRKOV, M. LYUTIKOV, L. BURZAWA

Radiotransients



Many different types of transient sources are already detected at radio wavelengths.

However, detection of very short and non-repeating flares of unknown sources without identification at other bands is a very complicated task.


Rotating Radio Transients (RRATs) – millisecond radio bursts from neutron stars, - have been identified in 2006.

In 2007 the first example of a new class of millisecond radio transients have been announced: the first extragalactic millisecond radio burst.

History of FRBs


- 2007 Lorimer et al. The first event announced.
- 2012 Keane et al. The second event.
- 2013 Thornton et al. Four events. The story really starts.
- 2016 Spitler et al. The first repeating source.
Chatterjee et al. Identification of the host galaxy

Lorimer et al.



Large dispersion measure points to extragalactic origin.

This is supported by isotropic sky distribution and many other considerations.



Catalogue


22 FRBs
+
one repeater

Rate: several thousands
per day per sky


Event	Telescope	gl [deg]	gb [deg]	FWHM [deg]	DM [cm ⁻³ pc]	S/N	W _{obs} [ms]	S _{peak,obs} [Jy]	F _{obs} [Jy ms]
FRB010125	parkes	356.641	-20.020	0.25	790(3)	17	9.40 ^{+0.20} _{-0.20}	0.30	2.82
FRB010621	parkes	25.433	-4.003	0.25	745(10)		7.00	0.41	2.87
FRB010724	parkes	300.653	-41.805	0.25	375	23	5.00	>30.00 ^{+10.00} _{-10.00}	>150.00
FRB090625	parkes	226.443	-60.030	0.25	899.55(1)	30	1.92 ^{+0.83} _{-0.77}	1.14 ^{+0.42} _{-0.21}	2.19 ^{+2.10} _{-1.12}
FRB110220	parkes	50.828	-54.766	0.25	944.38(5)	49	5.60 ^{+0.10} _{-0.10}	1.30 ^{+0.00} _{-0.00}	7.28 ^{+0.13} _{-0.13}
FRB110523	GBT	56.119	-37.819	0.26	623.30(6)	42	1.73 ^{+0.17} _{-0.17}	0.60	1.04
FRB110626	parkes	355.861	-41.752	0.25	723.0(3)	11	1.40	0.40	0.56
FRB110703	parkes	80.997	-59.019	0.25	1103.6(7)	16	4.30	0.50	2.15
FRB120127	parkes	49.287	-66.203	0.25	553.3(3)	11	1.10	0.50	0.55
FRB121002	parkes	308.219	-26.264	0.25	1629.18(2)	16	5.44 ^{+3.50} _{-1.20}	0.43 ^{+0.33} _{-0.06}	2.34 ^{+4.46} _{-0.77}
FRB121102	arecibo	174.950	-0.225	0.05	557(2)	14	3.00 ^{+0.50} _{-0.50}	0.40 ^{+0.40} _{-0.10}	1.20 ^{+1.60} _{-0.45}
FRB130626	parkes	7.450	27.420	0.25	952.4(1)	21	1.98 ^{+1.20} _{-0.44}	0.74 ^{+0.49} _{-0.11}	1.47 ^{+2.45} _{-0.50}
FRB130628	parkes	225.955	30.655	0.25	469.88(1)	29	0.64 ^{+0.13} _{-0.13}	1.91 ^{+0.29} _{-0.23}	1.22 ^{+0.47} _{-0.37}
FRB130729	parkes	324.787	54.744	0.25	861(2)	14	15.61 ^{+9.98} _{-6.27}	0.22 ^{+0.17} _{-0.05}	3.43 ^{+6.55} _{-1.81}
FRB131104	parkes	260.549	-21.925	0.25	779(1)	30	2.08	1.12	2.33
FRB140514	parkes	50.841	-54.611	0.25	562.7(6)	16	2.80 ^{+3.50} _{-0.70}	0.47 ^{+0.11} _{-0.08}	1.32 ^{+2.34} _{-0.50}
FRB150215	parkes	24.662	5.280	0.25	1105.6(8)	19	2.80 ^{+1.20} _{-0.50}	0.70 ^{+0.28} _{-0.01}	1.96 ^{+1.96} _{-0.37}
FRB150418	parkes	232.665	-3.234	0.25	776.2(5)	39	0.83 ^{+0.25} _{-0.25}	2.19 ^{+0.60} _{-0.30}	1.82 ^{+1.20} _{-0.72}
FRB150807	parkes	336.709	-54.400	0.25	266.5(1)		0.35 ^{+0.05} _{-0.05}	128.00 ^{+5.00} _{-5.00}	44.80 ^{+8.40} _{-7.90}
FRB160317	UTMOST	246.050	-0.990	0.00	1165(11)	13	21.00 ^{+7.00} _{-7.00}	>3.00	>63.00
FRB160410	UTMOST	220.360	27.190	0.00	278(3)	13	4.00 ^{+1.00} _{-1.00}	>7.00	>28.00
FRB160608	UTMOST	254.110	-9.539	0.00	682(7)	12	9.00 ^{+6.00} _{-6.00}	>4.30	>38.70
FRB170107	ASKAP	266.000	51.400	0.13	609.5(5)	16	2.60	22.30	57.98

Localization


Radius of uncertainty circle ~ 10 arcmin



Usually FRBs are seen just in one beam.



Statistical properties of FRBs



$$\frac{dN}{dF_{\text{obs}}} = (4.4 \pm 0.4) \times 10^3 F_{\text{obs}}^{-1.18 \pm 0.15} \text{ sky}^{-1} \text{ day}^{-1}$$

FRBs. Different hypotheses

Millisecond extragalactic radio bursts of that intensity without immediate identification with other bursts have not been predicted by earlier studies.

Since 2007 many hypotheses have been proposed.

A real flow started in late summer of 2013 after the paper by Thornton et al.

- Magnetars
- Super radio pulsars
- Evaporating black holes
- Coalescing NSs
- Coalescing WDs
- Coalescing NS+BH
- Supramassive NSs
- Deconfinement of a NS
- Axion clouds and NSs
- Cosmic strings
- Charged BHs
- NS collapse




Magnetar model

The first idea of possible connection between FRBs and magnetars has been proposed already in 2007 by Popov, Postnov: arXiv 0710.2006.


This hypothesis has been based on rate and energetics considerations, mainly. FRB bursts might be related to giant flares of magnetars

Later this approach was developed by Lyubarsky (2014).

In the model by Lyubarsky the radio burst happens due to synchrotron maser emission after interaction between a magnetic pulse after a giant flare of a magnetar with surrounding nebula.




Nebula emission




The model of a nebular emission after a huge energy release in a central source was developed by several authors.


1603.08875



Magnetar






RNS





MWD

Nebulae around magnetars

1206.3330. New results in 1604.06472



There are examples of nebulae around magnetars and highly magnetized radio pulsars.



0909.3843


About formation of pulsar nebulae around magnetars see 1606.01391

1211.0852


Radio flares from M31

Rubio-Herrera et al. (2013)
discovered millisecond radio bursts
from the Andromeda galaxy.

It looks like a scaled version of FRBs.
In the magnetar model such (more frequent) bursts
can be related to weaker flares of magnetars.




Note, that Frederiks et al. (2005)
proposed a candidate for a
giant magnetar flare in M31.



Radio pulsar model

1004.2803



In the case of the Crab pulsar so-called giant pulses are known.

It has been suggested (1501.00753, 1505.05535) that young pulsars with large \dot{E} can rarely produce much more energetic events.

Scaling allows to reproduce energetics of FRBs.

FRBs as supergiant pulses

$$\eta = \frac{L_{GP}}{\dot{E}_{Crab}} = \frac{\nu c^3 d_{Crab}^2 S_\nu P_{NS}^4}{4\pi^3 B_{NS}^2 R_{NS}^6} \approx 10^{-2},$$

Estimates are done via scaling of parameters of the Crab. Rather normal magnetic field but rapid rotation formally can explain FRB energetics.

$$L_{FRB} = \eta \dot{E} \rightarrow B_{NS} = \frac{c^{3/2} d \sqrt{(\nu F_\nu)} P_{NS}^2}{2\pi^{3/2} R_{NS}^{3/2} \sqrt{\eta}} = 2 \times 10^{13} d_{100\text{Mpc}} F_{30\text{Jy}}^{1/2} \tau_{5\text{msec}}^2 \sqrt{\nu_9 \eta_{-2}^{-1/2}} \text{ G.}$$

$$\tau_{SD} = \frac{\pi \eta I_{NS}}{d^2 F_\nu \mu P^2} \sim \text{few years.}$$

With magnetic field and spin period it is possible to estimate the characteristic spin-down time.

Supergiant pulses of young radio pulsars in dense supernova remnants

Age 30-100 years

Uniform distribution in \dot{E} in logarithmic scale

Absorption of low-frequency radiation in the remnant


Repeating bursts

Bursts are uniformly distributed in distance

$$f(\dot{E}) \propto \frac{\ln(\dot{E}_0/\dot{E})}{\dot{E}^{3/2}}, \beta = 1$$

Steady state solution for magneto-dipole spindown and

$$f_{inj}(\dot{E}) \propto \dot{E}^{-\beta}$$



Dispersion in a dense supernova remnant

$$\text{DM} \approx \frac{M_{ej}}{m_p r^2}$$

$$r = \sqrt{M_{ej}/m_p} \frac{1}{\sqrt{\text{DM}}} = 0.34 \text{pc} \sqrt{m_\odot} \text{DM}_{375}^{-1/2}$$

Dispersion in a dense SNR might explain observed DM of FRBs in the model when they are near-by at distances $\sim 100\text{-}200$ Mpc.

$$\frac{M_{swept}}{M_{ej}} = \sqrt{M_{ej}/m_p} \frac{n_{ISM}}{\text{DM}^{3/2} \text{pc}^{3/2}} = 4.5 \times 10^{-4} n_{ISM} \sqrt{m_\odot} \ll 1,$$

$$v_{ej} = \sqrt{\frac{2E_{ej}}{M_{ej}}}$$

$$t = \frac{M_{ej}}{\sqrt{2\text{DM}E_{ej}m_p}} = 35 \text{yrs} m_\odot$$

$$\tau = 8 \times 10^{-2} n^2 \nu^{-2.1} r T^{-1.35} = 0.05 \text{DM}_{375}^{5/2} m_\odot^{-1/2} \nu_9^{-2.1}$$

Burst rate

SN rate $\sim 3 \cdot 10^{-4} \text{ yr}^{-1} \text{ Mpc}^{-3}$ (Dahlen et al. 2012).

This gives ~ 1 SN per day in 100 Mpc.

Ages and typical lifetime of our sources ~ 30 -100 years.

Thus, we have $\sim 10\,000 - 30\,000$ sources in 100 Mpc.

The observed rate of FRBs $\sim 3 \cdot 10^3$ per day.

Then, each source might give a flare per few days.

If we increase the distance up to 200 Mpc then we can use just 10% of most energetic neutron stars.

Giant pulses of the Crab with fluence 100-200 kJy for Edot increase by factor 100 000 are scaled to flares with the flux ~ 1 Jy from 100-200 Mpc.

Number of giant pulsars depends on flux as $\sim S^{-3}$.


For FRBs we then obtain that most bright event might be observed once per few months.

Monte Carlo simulations


$$f_{inj} \propto \dot{E}^{-1}$$

$$\partial_t \dot{E} \propto -\dot{E}^{-3/2}$$

$$S \propto \dot{E}/r^2.$$



Distribution in distance and DM



$$S \propto \frac{\dot{E}}{r^2} \propto \dot{E} f(\dot{E})^{2/3}$$

$$S \propto \dot{E}^{(1-2\alpha/3)} \propto r^{-2+3/\alpha}$$

$$\dot{E} = \frac{\dot{E}_0}{(1 + t/\tau)^2},$$

$$DM = \frac{M_{ej}^2}{2E_{ej}m_p\tau} \frac{\dot{E}}{(\dot{E} + \dot{E}_0)^2}$$

Dispersion measure does not correlate with fluence or peak luminosity.

This is in correspondence with the model.

FRB vs. ULX

For a typical FRB with peak flux $S_{\text{peak}} = 1$ Jy we obtain radio luminosity:

$$L_r = 1.7 \times 10^{40} (S_{\text{peak}}/1 \text{ Jy}) (d/100 \text{ Mpc})^2 \text{ erg s}^{-1}.$$

Then, rotational energy losses are:

$$\dot{E} = 1.7 \times 10^{42} (S_{\text{peak}}/1 \text{ Jy}) (d/100 \text{ Mpc})^2 (\eta/0.01)^{-1} \text{ erg s}^{-1}.$$

Using the relation from Possenti et al. we obtain the X-ray luminosity:

$$L_X = 1.8 \times 10^{41} (S_{\text{peak}}/1 \text{ Jy})^{1.34} \times \\ \times (d/100 \text{ Mpc})^{2.68} (\eta/0.01)^{-1.34} \text{ erg s}^{-1}.$$

And so, the X-ray flux is:

$$f_X = 1.5 \times 10^{-13} (S_{\text{peak}}/1 \text{ Jy})^{1.34} \times \\ \times (d/100 \text{ Mpc})^{0.68} (\eta/0.01)^{-1.34} \text{ erg cm}^{-2} \text{ s}^{-1}.$$

For large distances we obtain higher f_X for a given S_{peak} , for smaller — weaker. If a source with peak flux 1 Jy is at 10 Mpc, then $f_X = 3.2 \times 10^{-14} \text{ erg cm}^{-2} \text{ s}^{-1}$. Correspondently, for 200 Mpc we have $f_X = 2.5 \times 10^{-13} \text{ erg cm}^{-2} \text{ s}^{-1}$.

In the model of supergiant pulses it is natural to expect that at distances 100-200 Mpc young energetic PSRs might be strong X-ray sources, similar to ULXs.

$$L_X \approx 2 \times 10^{42} \left(\dot{E}/10^{43} \text{ erg s}^{-1} \right)^{1.34} \text{ erg s}^{-1},$$

(Possenti et al. 2002)

Searches for possible counterparts of FRBs in X-ray in near-by (100-200 Mpc) galaxies can confirm or falsify the model.

Future observations

FAST



FAST – burst per week

1602.06099



SKA

SKA – burst per hour!

1602.05165, 1501.07535


Special projects partly dedicated to FRBs



<https://sites.google.com/site/publicsuperb/>



1607.02059



CHIME – burst per day!
1601.02444

HIRAX.
South variant of CHIME


<http://chime.phas.ubc.ca/>

ASKAP and Apertif

ASKAP



Few bursts per week.
1709.02189



Westerbork

Northern sky.
Doubling the number?
Rapid on-line
identification – follow-up

